

UNIVERSITY OF LIÈGE

FACULTY OF SCIENCES

DEPARTMENT OF GEOGRAPHY

**Representation of Foehn events in the
Modèle Atmosphérique Régional (MAR)
in Patagonia**

Can a change in DEM generalization improve the model?

Author

Adrien DAMSEAUX

Supervisors

Dr. X. FETTWEIS

Pr. Y. CORNET

Jury president

Dr. X. FETTWEIS

Reader

Dr. G. HOUBRECHTS

ERRATA

This thesis is submitted for the degree of
MASTER IN GEOGRAPHICAL SCIENCES, CLIMATOLOGY ORIENTATION

Academic year 2017-2018

Defence date: September 2018




Figure 1 Four profiles of the vertical surface elevation. The aim is to show the performances of each DEM generalization method with respect to the original DEM. Background: DEM ETOPO1 at a 1 arc-minute resolution.




Figure 2 Surface elevation profile A. Resolution: 1 arc-minute for ETOPO1, 10 km for the others.




Figure 3 Surface elevation profile B. Resolution: 1 arc-minute for ETOPO1, 10 km for the others.




Figure 4 Surface elevation profile C. Resolution: 1 arc-minute for ETOPO1, 10 km for the others.




Figure 5 Surface elevation profile D. Resolution: 1 arc-minute for ETPO1, 10 km for the others.